

RESEARCH NOTE

Megaselia scalaris* (Loew) (DIPTERA: PHORIDAE) INFESTING EDIBLE WILD MUSHROOM, *Boletus griseipurpureus

LAU, M.F., ROSNIDA, T., SUHAILA, A.B. and LATIFFAH, Z.*

School of Biological Sciences, Universiti Sains Malaysia, 11800 USM Penang, Malaysia

*Email: Lfah@usm.my / latiffahz@yahoo.com

Accepted 29 May 2017, Published online 27 June 2017

The genus *Megaselia* Rondani of the family Phoridae (Diptera) is the most numerous phorid species and distributed in every geographic region (Disney, 2008). One of the phorid species known as scuttle fly, *Megaselia scalaris* (Loew) is cosmopolitan and synanthropic species (Costa *et al.*, 2007). The larvae of *M. scalaris* develop in various types of habitats, consuming a wide range of organic materials from plant and animal remains than any other insect known. Thus, the larvae is often described as detritivore, parasite, facultative parasite, and parasitoid (Tumrasvin *et al.*, 1997; Disney, 2008, Mongiardino Koch *et al.*, 2013). As *M. scalaris* can occupied a wide range of habitat and has spectrum of feeding behaviour, this scuttle fly has been reported in many instances including in forensic cases (Varney & Noor, 2010), myiasis in humans and animals (Disney, 2008; Ghavami & Djalilvand, 2015), as laboratory pest (Miller, 1979; Costa *et al.*, 2007; Garriss, 2014) and has been used in genetic, developmental and bioassay studies (Disney, 2008).

Among the species that *Megaselia* have been reported as pest of cultivated and non-cultivated mushroom, *M. halterata*, is the most common and widespread pest of cultivated mushroom (Civelek & Önder, 1997; Lewandowski *et al.*, 2012) having infesting cultivated white button mushroom (*Agaricus bisporus*) (Erler and Polat, 2015) while *M. tamilnaduensis* infested oyster mushrooms (*Pleurotus* spp.) (Mohan *et al.*, 1996). The fruiting bodies of *Termitomyces albuminosus* (non-cultivated mushroom) was attacked by *M. termitomycana* (Disney & Chou 1996), *Pulveroboletus ravenelii* by *M. pulveroboleti* (Disney & Chou, 1998) and *Rigidoporus microporus* by *M. sororbata* (Disney & Ševčík, 2011). In Japan, fruiting bodies of fungi from the

genera *Russula* and *Gymnopilus* were reported to be infested by *M. donaldsonae*, *M. flava*, *M. gotoi*, *M. kanekoi*, *M. margaretae*, *M. nakayamai*, *M. salteri* and *M. stepheni* (Disney *et al.*, 2014). Based on reports by Disney (1994) and Ševčík (2001), only three species of *Megaselia*; *M. lata*, *M. flavicans*, *M. berndseni* (synonym: *M. pygmaeoides*) reported by Bruns (1984) have been recovered from *Boletus* mushrooms.

However, to our knowledge, there is no report on the infestation of *Megaselia* on mushroom in Malaysia. The present study reported the first record of *M. scalaris* infesting wild mushroom, *Boletus griseipurpureus* found in peat swamp forest in Bachok, Kelantan, Peninsular Malaysia.

The wild mushroom, *Boletus griseipurpureus* were collected from swamp forest or marshland, 7 m above mean sea level with latitude 05° 56' 00" and longitude 102° 25' 00" in the district of Bachok, Kelantan, Peninsular Malaysia. A total of 96 fruiting bodies were collected from the sampling sites during fruiting season from late June 2011 to end of September 2012 before the northeast monsoon commenced along the east coast of Peninsular Malaysia.

The *B. griseipurpureus* fruiting bodies harbouring the larvae were transferred into a 500 ml conical flask, covered with double layered of paper towel and secured by a rubber band. The larvae were reared to the adult stage in an incubator at 30°C. The adult flies were collected, mounted on microscopic slides and identified. Identification of the pest was performed based on taxonomic keys and descriptions of Disney (1994), Borror & White (1998) and Triplehorn *et al.* (2005) by using a digital camera (Olympus, Xcam-α) attached to a microscope (Olympus, Model BX41).

From 96 fruiting bodies collected, 54 samples (about 60%) were infested by the larvae that were found in the pore tube and the pileal context (Figures

* To whom correspondence should be addressed.




Fig. 1. Infestation by the insect larvae. (A) In the pore tube. (B) In the pileal context.




Fig. 2. Antennal structure of male *Megaselia scalaris*. (A) Globular third antennal segment. (B) Apical pseudo-arista.

1A and 1B). Some eggs embedding in the pores were also observed. After two weeks, three adult flies were collected from the rearing. The flies were identified as male *Megaselia scalaris* (Loew) (Diptera: Phoridae) (Figures 2A and 2B) and was confirmed by Dr. Brian V. Brown (Curator, Entomology Section, National History Museum of Los Angeles County, Los Angeles, USA).

Megaselia attacks a wide range of macrofungi such as *Auricularia* sp., *Pleurotus* sp., *Rigidoporus* sp. and *Termitomyces* sp. as they are attracted to putrid odors from the mushrooms (Diyes *et al.*, 2015). Yamashita *et al.* (2005) reported three species of scuttle fly, namely *M. flava*, *M. kanekoi* and *M. gotoi* feed on deadly and poisonous *Amanita ibotengutake*; and suggested that the edibility of *B. griseipurpureus* could not be drawn from the

presence of *M. scalaris* larvae. Besides mushroom structure and size (Yamashita & Hijii, 2003), the infestation by *Megaselia* is also attributed to other factors, including host species (Smith *et al.*, 2006), type of food sources (Idris *et al.*, 2001), abundance of fungal resources (Takahashi *et al.*, 2005) and ambient temperature (Barzegar *et al.*, 2011). It was believed that *B. griseipurpureus* has the potential to be the selective host for *M. scalaris* in peat swamp forest. Limiting fluid sugary meals in the swamp forest may in particular make the *M. scalaris* to choose *B. griseipurpureus* as a host species. According to Sukontason *et al.* (2006), *M. scalaris* access fluid meals primarily on sugar because it has sponging mouthparts and Kurunaweera *et al.* (2002) also found *M. scalaris* on flesh ripe banana, which contained high sugar level.

Mushrooms are high in protein, fiber, and carbohydrates (Cheung, 2010) suggesting that *M. scalaris* utilized protein sources from *B. griseipurpureus*. Protein concentration in *B. griseipurpureus* is 34% (Gbolagade *et al.*, 2006) while general crude protein of edible mushrooms range from 15-35% (Cheung, 2010). *Boletus griseipurpureus* with its large fruiting body could provide sufficient amount of protein source for larval development of *M. scalaris*. Corresponding to the current observation, *B. griseipurpureus* is associated with the same host tree and thus produce fruiting bodies at the same site annually in Thailand (Seehanan & Petcharat, 2008; Aung-Aud-Chariya *et al.*, 2012). The mushroom abundance and predictable fruiting bodies location might account for the infestation of *M. scalaris* in the peat swamp forests. Indeed, Diyes *et al.* (2015) mentioned *M. scalaris* preferred to lay their eggs on decaying plant matter (in this case *B. griseipurpureus* live on them). The fine structure of eggs (boat-shaped with gunwale-like palisade of flat platelets form) (Disney, 1994; Harrison & Cooper, 2003; Disney, 2008) fit well in the tubes indicating that the pileus of *B. griseipurpureus* provided an ideal breeding site for *M. scalaris*. Although no visible damage of the fruiting body was observed, the spores might fail to disperse as a consequence of the blockage by the embedded eggs. The decaying plant was greatly found in peat swamp forest do encourage the abundance of mushroom resources. Furthermore, *M. scalaris* larvae is depending on moist from the decaying plant material to survive (Sukontason *et al.*, 2006).

In conclusion, wild *B. griseipurpureus* can be the potential selective host for *M. scalaris* in peat swamp forests and reveals the possibility of *M. scalaris* infesting the edible wild mushroom, *B. griseipurpureus* in Malaysia.

ACKNOWLEDGEMENTS

The authors wish to thank Dr. Brain V. Brown (Curator, Entomology Section, National History Museum of Los Angeles County, Los Angeles, USA) for the fly species confirmation and his valuable comments. This study was funded in part by Research University Grant 1001/PBIOLOGY/811179.

REFERENCES

Aung-aud-chariya, A., Bangrak, P., Dell, B., Lumyong, S. & Kamlangdee, N. 2012. Preliminary molecular identification of *Boletus griseipurpureus* (Corner) from Thailand and

its nutritional value. *Journal of Agricultural Technology*, **8**: 1991-1998.

Barzegar, S., Zamani, A.A., Abbasi, S. & Shooshtari, R.V. 2011. Effect of temperatures on life expectancy and reproductive parameters of *Megaselia halterata* (Diptera: Phoridae) on different varieties of button mushroom. *Euphrates Journal of Agriculture Science*, **3**: 147-151.

Borror, D.J. & White, R.E. 1998. A field guide to insects: America North of Mexico. Houghton Mifflin Company, Boston, New York.

Bruns, T.D. 1984. Insect mycophagy in the Boletales: Fungivore diversity and the mushroom habitat. In: Fungus-insect relationships: Perspectives in ecology and evolution. Wheeler, Q and Blackwell, M. (eds.). Columbia University Press. pp. 91-129.

Cheung, P.C.K. 2010. The nutritional and health benefits of mushrooms. *Nutrition Bulletin*, **35**: 292-299.

Civelek, H.S. & Önder, F. 1997. The harmful Diptera species in mushroom produce stores in İzmir province. Proceedings of the Third Turkish National Congress of Entomology, 24-28 September 1996, Ankara, University of Ankara Press, pp. 716.

Costa, J., Almeida, C.E.G., Esperança, M., Morales, N., Jacenir, R. dos S. Mallet, Teresa, C.M. & Gonçalves, A.P. do Prado. 2007. First record of *Megaselia scalaris* (Loew) (Diptera: Phoridae) infesting laboratory colonies of *Triatoma brasiliensis neiva* (Hemiptera: Reduviidae). *Neotropical Entomology*, **36**(6): 987-989.

Disney, R.H.L. 1994. Scuttle flies: The Phoridae. Chapman and Hall, London.

Disney, R.H.L. 2008. Natural history of the scuttle fly, *Megaselia scalaris*. *Annual Review of Entomology*, **53**: 39-60.

Disney, R.H.L. & Chou, W.N. 1996. A new species of *Megaselia* (Diptera: Phoridae) reared from the fungus *Termitomyces* (Agaricales: Amanitaceae) in Taiwan. *Zoological Studies*, **35**: 215-219.

Disney, R.H.L. & Chou, W.N. 1998. A new species of *Megaselia* (Diptera: Phoridae) reared from the fungus *Pulveroboletus* (Boletales: Boletaceae) in Taiwan. *Bulletin of the National Museum of Natural Science, Taiwan*, **11**: 135-139.

Disney, R.H.L. & Ševčík, J. 2011. A new species of fungus breeding *Megaselia* (Diptera: Phoridae) from Indonesia. *Acta Entomologica Musei Nationalis Pragae*, **51**: 211-215.

Disney, R., Henry, L., Masayuki, N., Mio, K. & Nobuko, T. 2014. New records of *Megaselia* (Diptera: Phoridae) reared from fungus sporophores in Japan, including five new species. *Applied Entomology and Zoology*, **49**(4): 541-552.

- Diyes, G.C.P., Karunaratne, W.A.I.P., Tomberlin, J.K. & Rajakaruna, R.S. 2015. First record of *Megaselia scalaris* (LOEW) (Diptera: Phoridae) infesting a spinose ear tick, *Otobius megnini*, colony in Sri Lanka. *Tropical Biomedicine*, **32**(4): 791-795.
- Erlar, F. & Polat, R. 2015. The flies on mushrooms cultivated in the Antalya-Korkuteli district and their control. *Akdeniz Üniversitesi Ziraat Fakültesi Dergisi*, **28**(2): 61-66.
- Garris, G.I. 2014. *Megaselia scalaris* (Diptera: Phoridae) infesting laboratory tick colonies. *Journal of Medical Entomology*, **20**: 688.
- Gbolagade, J., Ajayi, A., Oku, I. & Wankasi, D. 2006. Nutritive value of common wild edible mushrooms from southern Nigeria. *Global Journal of Biotechnology and Biochemistry*, **1**: 16-21.
- Ghavami, M.B. & Djalilvand, A. 2015. First record of urogenital myiasis induced by *Megaselia scalaris* (Diptera: Phoridae) from Iran. *Journal of Arthropod Borne Disease*, **9**(2): 274-280.
- Harrison, D.A. & Cooper, R.L. 2003. Characterization of development, behavior and neuromuscular physiology in the phorid fly, *Megaselia scalaris*. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology*, **136**: 427-439.
- Idris, A.B., Abdullah, M. & Lin, T.P. 2001. Effect of various diets on the development of scuttle fly, *Megaselia scalaris* (Loew) (Diptera: Phoridae), larvae and pupae and percent of adult emergence and longevity. *Pakistan Journal of Biological Sciences*, **4**: 321-325.
- Karunaweera, N.D., Ithalamulla, R.L. & Kumarasinghe, S.P. 2000. *Megaselia scalaris* (Diptera: Phoridae) can live on ripe bananas – a potential health hazard? *Ceylon Medical Journal*, **47**: 9-10.
- Lewandowski, M., Kozak, M. & Sznyk-basałyga, A. 2012. Biology and morphometry of *Megaselia halterata*, an important insect pest of mushrooms. *Bulletin of Insectology*, **65**: 1-8.
- Miller, P.L. 1979. Intraspecific phorid phoresy. *Entomologist's Monthly Magazine*, **114**: 211-214.
- Mohan, S., Mohan, S. & Disney, R.H.L. 1996. A new species of scuttle fly (Diptera: Phoridae) that is a pest of oyster mushrooms (Agaricales: Pleurotaceae) in India. *Bulletin of Entomological Research*, **85**: 515-518.
- Mongiardinio Koch, N., Fontanarrosa, P., Padró, J. & Soto, M. 2013. First record of *Megaselia scalaris* (Loew) (Diptera: Phoridae) infesting laboratory stocks of mantids (*Parastagma-toptera tessellata*, Saussure). *Arthropods*, **2**(1): 1-6.
- Seehanan, S. & Petcharat, V. 2008. Some species of wild Boletes in Thailand. *Journal of Agriculture Technology*, **4**: 109-118.
- Ševčík, J. 2001. Diptera (excluding Mycetophilidae S. str.) associated with fungi in Czech and Slovak Republics: A survey of rearing records from 1998-2000. *Acta Universitatis Carolinae Biologica*, **45**: 157-168.
- Smith, J.E., Challen, M.P., White, P.F., Edmondson, R.N. & Chandler, D. 2006. Differential effect of *Agaricus* host species on the population development of *Megaselia halterata* (Diptera: Phoridae). *Bulletin of Entomological Research*, **96**: 565-571.
- Sukontason, K.L., Boonsriwong, W., Siri wattana-rungsee, S., Piangjai, S. & Sukontason, K. 2006. Morphology of puparia of *Megaselia scalaris* (Diptera: Phoridae), a fly species of medical and forensic importance. *Parasitology Research*, **98**: 268-272.
- Takahashi, K.H., Tuno, N. & Kagaya, T. 2005. Abundance of mycophagous arthropods present on different species of fungi in relation to resource abundance at different spatial scales. *European Journal of Entomology*, **102**: 309-46.
- Triplehorn, C.A., Johnson, N.F. & Borror, D.J. 2005. Borror and DeLong's introduction to the study of insects. Seventh edition. Thompson Brooks/Cole, University of Minnesota.
- Tumrasvin, W., Sucharit, S. & Vutikes, S. 1997. Studies on the life history of *Megaselia scalaris* (Loew) in Thailand. *Southeast Asian Journal of Tropical Medicine and Public Health*, **8**: 74-76.
- Yamashita, H., Tanaka, C., Nakayama, H., Tuno, N. & Osawa, N. 2005. New host record for three scuttle flies, *Megaselia flava*, *M. kanekoi* and *M. gotoi* (Diptera: Phoridae), on the poisonous fungus *Amanita ibotengutake* (Agaricales: Amanitaceae). *Entomological Sciences*, **8**: 223-225.
- Yamashita, S. & Hijii, N. 2003. Effects of mushroom size on the structure of a mycophagous arthropod community: Comparison between infracommunities with different types of resource utilization. *Ecological Research*, **18**: 131-143.
- Varney, R.L. & Noor, M.A.F. 2010. The scuttle fly. *Current Biology*, **20**(11): 466-467.